arXiv:0903.4459v1 [math.AG] 25 Mar 2009

MORPHISMS OF COHOMOLOGICAL FIELD THEORIES

KHOA LU NGUYEN AND CHRIS WOODWARD

Abstract. We introduce a notion of morphism of CohFT's, on the basis of the analogy with A_∞ morphisms, and discuss the relationship with morphisms of F-manifolds introduced by Manin and Hertling [5]. The structure maps of a morphism of CohFT's have as input a cohomology class on the moduli space of *scaled affine lines* (complexified multiplihedron) studied in Ma'u-Woodward [9]. The main result is a computation of the space of Cartier divisors on this moduli space; these appear in the splitting axiom.

1. Introduction

In order to formalize the algebraic structure of Gromov-Witten theory Kontsevich and Manin introduced a notion of *cohomological field theory* (CohFT), see [8, Section IV]. The correlators of such a theory depend on the choice of cohomological classes on the moduli space of stable curves $\overline{M}_{g,n}$, and satisfy a splitting axiom for each boundary divisor of $\overline{M}_{g,n}$. In genus zero $\overline{M}_{0,n}$ may be viewed as the complexification of Stasheff's associahedron and the notion of CohFT is closely related to the notion of A_∞-algebra. Dubrovin constructed from any CohFT a *Frobenius manifold*, which is a manifold with a family of multiplications on its tangent spaces together with some additional data. Later Hertling-Manin [5] proposed the notion of *F-manifold*, which remembers only the multiplications.

In this paper we introduce a notion of *morphism* of CohFT's which is a "closed string" analog of a morphism of A_∞ -algebras. The additional data in the structure maps is the choice of cohomological classes on the *complexified multiplihedra* or *moduli of scaled curves* $\overline{M}_{n,1}(\mathbb{A})$, introduced in Ziltener [12]. This space was studied in Ma'u-Woodward [9] where it was shown that it may be viewed as the complexification of Stasheff's multiplihedron, which appears in the definition of A_∞ map. The splitting axiom for a morphism of CohFT's requires that for each Cartier divisor on $\overline{M}_{n,1}$ one has a splitting axiom similar to that for CohFT's themselves. Any morphism of CohFT's gives rise to a map of the corresponding Frobenius manifolds, whose differential at any tangent space is an algebra homomorphism. In the language of [5], this is a *morphism of F-manifolds*.

The definition of morphism of CohFT is motivated by an attempt to extend the mirror theorems of Givental and others beyond the semipositive case in a more systematic

---

K.N. was supported by Undergraduate Research Experience portion of NSF grant DMS060509.

C.W. was partially supported by NSF grant DMS060509.

way than has been done previously. In Woodward-Ziltener [11], it is proposed that a morphism of CohFT's relates the gauged Gromov-Witten invariants of a variety with group action and the Gromov-Witten invariants of its geometric invariant theory quotient for a fixed underlying curve. In other words, morphisms of CohFT's provide an "algebraic home" for the counts of "vortex bubbles" that first appeared in Gaio-Salamon [3]. Thus morphisms of CohFT's are proposed as generalizations of the "mirror maps" in Givental [4], who studied semipositive toric varieties and complete intersections therein. There have already been a number of attempts in this direction, see for example Iritani [6]. However, we do not discuss this direction here. Neither do we discuss the relation with higher genus invariants, which seems somewhat mysterious to us at the moment (although morphisms of CohFT's can be expected to intertwine higher genus correlators for a fixed curve [11]).

The contents of the paper are as follows. Section 2 describes the definition of morphisms of CohFT in more detail. In Sections 3- 5, we use the local toric description of $\overline{M}_{n,1}(\mathbb{A})$ in [9] we give an explicit description of which boundary divisors are Cartier; this makes the definition precise. Section 3 describes combinatorially the cones of the toric singularities that appear. Section 4 computes the space of Cartier divisors in these toric varieties. Section 5 describes which combinations of boundary divisors in $\overline{M}_{n,1}(\mathbb{A})$ are Cartier, and contains the main result of this note: the relations on the space of Cartier divisors are the kernel of the push-pull map defined by the incidence relation on subsets and partitions. The last Section 6 sketches a notion of *composition of morphisms of CohFT*'s. Essentially composition is not quite defined by rather depends on the choice of additional data. However, there is a good notion of a *commutative triangle* of CohFT's.

We thank Ezra Getzler, Sikimeti Ma'u, Joseph Shao, and Constantin Teleman for helpful discussion and comments.

2. The definition

Let $\overline{M}_n$ denote the Grothendieck-Knudsen moduli space of genus zero n-marked stable curves, a smooth projective variety of dimension $\dim(\overline{M}_n) = n-3$. The boundary of $\overline{M}_n$ consists of the following divisors: for each splitting $\{1,\ldots,n\} = I_1 \cup I_2$ with $|I_1|, |I_2| \geq 2$ a divisor

$$\iota_{I_1\cup I_2} : D_{I_1 \cup I_2} \to \overline{M}_n$$

corresponding to the formation of a separating node, splitting the surface into pieces with markings I_1, I_2. The divisor $D_{I_1\cup I_2}$ is isomorphic to $\overline{M}_{|I_1|+1} \times \overline{M}_{|I_2|+1}$. Let

$$\delta_{I_1\cup I_2} \in H^2(\overline{M}_n)$$

denote its dual cohomology class. For any $\beta \in H(\overline{M}_n)$, let

$$\iota^*_{I_1\cup I_2}\beta = \sum_j \beta_{1,j} \otimes \beta_{2,j} \tag{1}$$

denote the Kunneth decomposition of its restriction to $D_{I_1\cup I_2}$.

Definition 2.1. An *(even, genus zero) cohomological field theory* is a datum $(V, (\mu^n)_{n\geq 2})$ where V is a vector space and $(\mu^n)_{n\geq 2}$ is a collection of *composition maps*

$$\mu^n : V^{\otimes n} \otimes H(\overline{M}_{n+1}, \mathbb{Q}) \to V$$

such that each μ^n is invariant under the action of the symmetric group S_n on V^n and the maps $(\mu^n)_{n\geq 2}$ satisfy a *splitting axiom*: for each partition $I_1 \cup I_2 = \{1, \ldots, n\}$, without loss of generality with $0 \in I_2$,

$$\mu^n(\alpha_1, \ldots, \alpha_n; \beta \wedge \delta_{I_1 \cup I_2}) = \sum_j \mu^{n-|I|+1}(\mu^{|I|}(\alpha_i, i \in I_1, \beta_{1,j}), \alpha_i i \notin I_2; \beta_{2,j})$$

where $\beta_{1,j}, \beta_{2,j}$ are as in (1).

The collection of composition maps $(\mu^n)_{n\geq 2}$ (which are termed in Manin [8] Comm_∞-structures) may be viewed as "complex analogs" of the A_∞ -structure maps of Stasheff, in the sense that the relevant moduli spaces have been "complexified". The various relations on the divisors in $\overline{M}_n$ give rise to relations on the maps μ^n. In particular the relation $[D_{\{0,3\}\cup\{1,2\}}] = [D_{\{0,1\}\cup\{2,3\}}]$ in $H^2(\overline{M}_4)$ implies that $\mu^2 : V \otimes V \to V$ is associative.

The notion of *morphism* of CohFT's based on the geometry of the *complexified multiplihedron* $\overline{M}_{n,1}(\mathbb{A})$ introduced in Ziltener [12] and studied further in Ma'u-Woodward [9]. Let $\mathbb{A}$ denote an affine line over a field k.

Definition 2.2. A *scaling* on $\mathbb{A}$ is an element of the space $\Omega^1(\mathbb{A})^k$ of translationally invariant one-forms. A *marked scaled line* is an affine line, which may assume is $\mathbb{A}$, equipped with a scaling $\alpha \in \Omega^1(\mathbb{A})^k$ and a collection $z_1, \ldots, z_n \in \mathbb{A}$ of distinct points.

From now on, we take $k = \mathbb{C}$. The moduli space $M_{n,1}(\mathbb{A})$ of scaled n-marked lines admits a compactification by allowing nodal curves as follows:

Definition 2.3. A *nodal marked scaled affine line* consists of

(a) A projective nodal curve $\underline{\Sigma} = (\Sigma_0, \ldots, \Sigma_n)$ of arithmetic genus zero,
(b) A collection $\underline{z} = (z_0, \ldots, z_n)$ of distinct, smooth points, such that $z_0 \in \Sigma_0$;
(c) A (possibly zero or infinite) scaling on Σ_i', where Σ_i' denotes the affine line obtained by removing the node connecting Σ_i to the root vertex, or z_0 if $i = 0$;

satisfying the condition that on any path from the root vertex Σ_0, to the component containing a marking, there is exactly one *colored component* with finite scaling; the components before (resp. after) this component have infinite (resp. zero scaling). A nodal marked scaled affine line is *stable* if each component with finite scaling has at least two special points, and each component with degenerate scaling has at least three special points. The *combinatorial type* of a nodal scaled affine line is the *rooted colored tree* $\Gamma = (\mathrm{V}(\Gamma), \mathrm{E}(\Gamma))$ whose vertices are the components of $\underline{\Sigma}$, edges are the nodes and markings, and *colored vertices* $\mathrm{V}^{\mathrm{col}}(\Gamma) \subset \mathrm{V}(\Gamma)$ the vertices corresponding to components with non-degenerate scalings.

Let $M_{n,1,\Gamma}(\mathbb{A})$ resp. $\overline{M}_{n,1}(\mathbb{A})$ denote the moduli space of scaled n-marked affine lines of type Γ resp. the union over combinatorial types. By [9] these moduli spaces admit the structure of quasiprojective resp. projective variety of dimension

$$\dim(M_{n,1,\Gamma}(\mathbb{A})) = n - 2 - \#\,\mathrm{E}(\Gamma) + \#\,\mathrm{V}^{\mathrm{col}}(\Gamma), \quad \dim(\overline{M}_{n,1}(\mathbb{A})) = n - 1. \tag{2}$$

The space $\overline{M}_{n,1}(\mathbb{A})$ was first studied in Ziltener's thesis [12] in the context of gauged Gromov-Witten theory on the affine line.

In the first non-trivial case $n = 2$, there exists an isomorphism

$$\overline{M}_{2,1}(\mathbb{A}) \to \mathbb{P}^1, \quad [z_1, z_2] \mapsto z_1 - z_2$$

with two distinguished points given by nodal scaled affine lines, appearing in the limit where the two markings become infinitely close or far apart. More generally, for arbitrary n there exists for any choice of $\{i, j\} \subset \{1, \ldots, n\}$ of subset of order 2 a *forgetful morphism*

$$f_{i,j} : \overline{M}_{n,1}(\mathbb{A}) \to \overline{M}_{2,1}(\mathbb{A})$$

forgetting the markings other than i, j and collapsing all unstable components, and for any choice $\{i, j, k, l\} \subset \{0, \ldots, n\}$ of subset of order 3 a forgetful morphism

$$f_{i,j,k,l} : \overline{M}_{n,1}(\mathbb{A}) \to \overline{M}_4$$

given by forgetting the scaling and all markings except i, j, k, l, and collapsing all stable components. The product of forgetful morphisms of the type above defines an embedding into a product of projective lines.

The variety $\overline{M}_{n,1}(\mathbb{A})$ is not smooth, but rather has toric singularities, see Section 2. The *boundary divisors* are the closures of strata $M_{n,1,\Gamma}$ of codimension one. From the dimension formula (2) one sees that there are two types of boundary divisors. First, for any $I \subset \{1, \ldots, n\}$ with $|I| \geq 2$ we have a divisor

$$\iota_I : D_I \to \overline{M}_{n,1}(\mathbb{A})$$

corresponding to the formation of a single bubble containing the markings I. This divisor admits a gluing isomorphism

$$D_I \to \overline{M}_{|I|+1} \times \overline{M}_{n-|I|+1,1}(\mathbb{A}). \tag{3}$$

Call these divisors of *type* I. Second, for any partition $I_1 \cup \ldots \cup I_r$ of $\{1, \ldots, n\}$ of size at least two we have a divisor $D_{I_1 \cup \ldots \cup I_r}$ corresponding to the formation of r bubbles with markings $I_1, \ldots, I_r$, attached to a remaining component with infinite area form. This divisor admits a gluing isomorphism

$$D_{I_1 \cup \ldots \cup I_r} \cong \left(\prod_{i=1}^{r} \overline{M}_{|I_i|,1}(\mathbb{A}) \right) \times \overline{M}_{r+1}. \tag{4}$$

Call these *divisors of type* II. Note that the divisors of type I and type II roughly correspond to the terms in the definition of A_∞ functor.

Since $\overline{M}_{n,1}(\mathbb{A})$ is not smooth, not every Weil divisor is Cartier, in particular, every divisor does not admit a dual cohomology class of degree 2. That is, for a Weil divisor

$$D = \sum_I n_I [D_I] + \sum_{I_1 \cup \ldots \cup I_r = \{1,\ldots,n\}} n_{I_1,\ldots,I_r} [D_{I_1,\ldots,I_r}] \tag{5}$$

there may or may not exist a class $\delta \in H^2(\overline{M}_{n,1}(\mathbb{A}))$ that satisfies

$$\langle \beta, [D] \rangle = \langle \beta \wedge \delta, [\overline{M}_{n,1}(\mathbb{A})] \rangle.$$

Let $(V, (\mu_V^n)_{n \geq 2})$ and $(W, (\mu_W^n)_{n \geq 2})$ be cohomological field theories.

Definition 2.4. A *strong morphism of cohomological field theories* is a collection of maps

$$\phi^n : V^{\otimes n} \otimes H(\overline{M}_{n,1}(\mathbb{A}), \mathbb{Q}) \to W, \quad n \geq 1$$

such that for any Cartier divisor D of the form (5) with dual class $\delta \in H^2(\overline{M}_{n,1}(\mathbb{A}))$ we have

$$\begin{aligned} \phi^n(\alpha, \beta \wedge \delta) = & \sum_I n_I \phi^{n-|I|+1}(\mu_V^{|I|}(\alpha_i, i \in I; \cdot), \alpha_j, j \notin I; \cdot)(\iota_I^* \beta) \\ & + \sum_{I_1,\ldots,I_r} n_{I_1,\ldots,I_r} \mu_W^r(\phi^{|I_1|}(\alpha_i, i \in I_r; \cdot), \ldots, \phi^{|I_r|}(\alpha_i, i \in I_r; \cdot); \cdot)(\iota^*_{I_1 \cup \ldots \cup I_r} \beta) \end{aligned} \tag{6}$$

where $\cdot$ indicates insertion of the Kunneth components of $\iota_I^*\beta$, $\iota^*_{I_1,\ldots,I_r}\beta$, using the homeomorphisms (3), (4). A *weak morphism* of CohFT's is defined similarly but with the additional data of an element $\phi^0 \in W$ and the splitting axiom allowing arbitrary numbers of insertions of ϕ^0, that is,

$$\begin{aligned} \phi^n(\alpha, \beta \wedge \delta) = & \sum_I n_I \phi^{n-|I|+1}(\mu_V^{|I|}(\alpha_i, i \in I; \cdot), \alpha_j, j \notin I; \cdot)(\iota_I^* \beta) \\ & + \sum_{r \leq s, I_1 \cup \ldots \cup I_r} n_{I_1 \cup \ldots \cup I_r} \mu_W^s(\phi^{|I_1|}(\alpha_i, i \in I_r; \cdot), \ldots, \\ & \qquad \phi^{|I_r|}(\alpha_i, i \in I_r; \cdot), \phi^0, \ldots, \phi^0; \cdot)(\iota^*_{I_1 \cup \ldots \cup I_r} \beta)/(s-r)!. \end{aligned} \tag{7}$$

A morphism of CohFT's is either a weak or strong morphism of CohFT's.

Example 2.5. $\overline{M}_{2,1}(\mathbb{A}) \cong \mathbb{P}^1$ and so every Weil divisor is Cartier and any two prime Weil divisors are linearly equivalent. In particular, the equivalence $[D_{\{1,2\}}] = [D_{\{1\},\{2\}}]$ holds in $H^2(\overline{M}_{2,1}(\mathbb{A}), \mathbb{Q}) = \mathbb{Q}$. Hence if $\phi : (V, (\mu_V^n)_{n \geq 2}) \to (W, (\mu_W^n)_{n \geq 2})$ is a strong morphism of CohFT's then $\phi^1 : V \to W$ is a homomorphism, $\phi^1 \circ \mu_V^2 = \mu_W^2 \circ (\phi^1 \times \phi^1)$. If $(\phi^n)_{n \geq 0}$ is a weak morphism of CohFT's, we have instead that

$$(\phi^1 \circ \mu_V^2)(v_1, v_2) = \sum_{s \geq 2} \mu_W^s(\phi^1(v_1), \phi^1(v_2), \phi^0, \ldots, \phi^0)/(s-2)! \tag{8}$$

for all $v_1, v_2 \in V$.

Recall that the notion of CohFT may be reformulated as a Frobenius manifold structure of Dubrovin [1], consisting of a datum $(V, g, F, 1, e)$ of an affine manifold V, a metric

g on the tangent spaces, a potential $F : V \to k$ whose third derivatives provide the tangent spaces T_tV with associative multiplications, a unity vector field 1 and an Euler vector field e providing a grading.

Proposition 2.6. *(see* [8]*) An CohFT* $(V, (\mu^n)_{n\geq 2})$ *has a canonical Frobenius structure with potential defined by* $F(v) = \sum_n \frac{1}{n!}\mu^n(v, \dots, v)$. *Conversely, any Frobenius structure defines a CohFT.*

Later, a weaker notion of *F-manifold* was introduced by Manin and Hertling [5], which consists of a pair $(V, \circ)$ where $\circ$ is a family of multiplications on the tangent spaces T_tV satisfying a certain axiom. In other words, one forgets the data $g, 1, e$. If $(V, \circ)$ and $(W, \circ)$ are F-manifolds then a *morphism of F-manifolds* is a smooth map $\phi : V \to W$ whose derivative $D_t\phi$ is a morphism of algebras from T_tV to $T_{\phi(t)}W$ for all $t \in V$. In particular, $D_0\phi = \phi^1$ is an algebra homomorphism from T_0V to $T_{\phi^0}W$. The F(robenius) interpretation of morphism of CohFT is the following:

Proposition 2.7. *Any morphism of CohFT's* $(\phi^n)_{n\geq 0}$ *from* V *to* W *defines a morphism of F-manifolds via the formula* $\phi : V \to W, \quad v \mapsto \sum_{n\geq 0} \frac{1}{n!}\phi^n(v, \dots, v)$.

Proof. Consider the relation $D_{\{1,2\}} \sim D_{\{1\},\{2\}}$ in $\overline{M}_{2,1}(\mathbb{A})$. Its pull-back to $\overline{M}_{n,1}(\mathbb{A})$ is the relation $\sum_{I_1,\dots,I_r} D_{I_1,I_2,\dots,I_r} \sim \sum_I D_I$ where the first sum is over partitions $I_1, \dots, I_r$ with $1 \in I_1, 2 \in I_2$, and the second is over subsets $I \subset \{1, \dots, n\}$ with $\{1, 2\} \subset I$; this follows easily from the definition of the forgetful morphisms. We obtain from the splitting axiom the relation

$$\begin{aligned}\sum_{n,i_1,\dots,i_r} \left((n - \sum_{j=1}^{r} i_j)!(i_1 - 1)!(i_2 - 1)!\prod_{j=3}^{r} i_j! \right)^{-1} \mu_W^r(\phi^{i_1}(a, t, \dots, t; 1) \\ \phi^{i_2}(b, t, \dots, t; 1), \phi^{i_3}(t, \dots, t; 1), \dots, \phi^{i_r}(t, \dots, t; 1); 1) = \\ \sum_{n,i}((i-2)!(n-i)!)^{-1}\phi^{n-i+1}(\mu_V^i(a, b, t, \dots, t), t, \dots, t; 1)\end{aligned}$$

which gives that $D_t\phi$ is a homomorphism. $\square$

It would be interesting to characterize which morphisms of F-manifolds arise from morphisms of CohFT's. This would require a study of the cohomology ring of $\overline{M}_{n,1}(\mathbb{A})$ along the lines of Keel [7] for the moduli space of stable marked genus zero curves; this paper is essentially a partial study of the second cohomology group only. The most naive possibility would be a positive answer to the following analog of Keel's result:

Naive Conjecture 2.8. $H(\overline{M}_{n,1}(\mathbb{A}))$ *is generated by the classes of the Cartier boundary divisors (described below in Theorem 5.1) modulo the following relations:*

(a) *the images of* $D_{\{1\}\{2\}} - D_{\{1,2\}}$ *under the forgetful morphism* $f_{ij}^* : \overline{M}_{n,1}(\mathbb{A}) \to \overline{M}_{2,1}(\mathbb{A})$*, as* i, j *range over distinct elements of* $\{1, \dots, n\}$*, and*
(b) *the products* $D'D''$*, if* D' *and* D'' *are disjoint Cartier divisors.*

3. Local structure of the moduli space of scaled curves

In this section we review the local description of the moduli space of scaled curves $\overline{M}_{n,1}(\mathbb{A})$ given in Ma'u-Woodward [9].

Definition 3.1. A *colored tree* Γ is a finite connected tree consisting of a set of vertices $\mathrm{V}(\Gamma)$, a set of edges $\mathrm{E}(\Gamma)$, a distinguished semi-infinite edge $e_0 \in \mathrm{E}(\Gamma)$ (called the *root*), and a subset of *colored vertices* $\mathrm{V}^{\mathrm{col}}(\Gamma) \subset \mathrm{V}(\Gamma)$ such that any non-self-crossing path from the root to any other semiinfinite edge crosses exactly one colored vertex. Γ is *stable* if the valence of any colored resp. uncolored vertex is at least two resp. three. The endpoint adjacent to the root edge is the *principal vertex* of Γ. Denote by $v_0, ..., v_{g-1}$ the uncolored vertices, with v_0 as the principal vertex, that is, the vertex adjacent to the root edge. Given two vertices v_i and v_j, we define v_j to be *below* v_i if there exists a non self-intersecting path starting from v_0 to some colored vertex such that the path passes through v_i before it does through v_j. We can define analogously this partial relation between two edges and between a vertex and an edge. Define the *principal branches* of a vertex v_k to be the edges below v_k which have v_k as one of their endpoints. The branches $d_1, ..., d_m$ of the principal vertex v_0 are called the *principal branches* of Γ, and the set of principal branches is denoted $\mathrm{B}(\Gamma)$

A map $\varphi : \mathrm{E}(\Gamma) \to \mathbb{C}$ is *balanced* if it satisfies the following condition: denote by $\mathrm{V}^-(\Gamma)$ the set of vertices on the root side of the colored vertices, that is, connected to the root by a path not crossing a colored vertex. For each vertex $v \in \mathrm{V}^-(\Gamma)$ and any colored vertex v' connected by a path of edges not crossing the root, let $p(v, v')$ denote the product of the values of φ along the unique path of edges from v to v'. Then φ is balanced if $p(v, v')$ is independent of the choice of colored vertex v'. Let $X(\Gamma)$ denote the set of balanced labellings:

$$X(\Gamma) := \{\varphi : \mathrm{E}(\Gamma) \to \mathbb{C} \,|\, \forall v \in \mathrm{V}^-(\Gamma),\ p(v, v') \text{ is independent of } v' \in \mathrm{V}^{\mathrm{col}}(\Gamma)\}. \tag{9}$$

The subset

$$T(\Gamma) = X(\Gamma) \cap \mathrm{Hom}(\mathrm{E}(\Gamma), \mathbb{C}^*)$$

of points with non-zero labels is the kernel of the homomorphism $\mathrm{Hom}(\mathrm{E}(\Gamma), \mathbb{C}^*) \to \mathrm{Hom}(\mathrm{V}^-(\Gamma), \mathbb{C}^*)$ given by taking the product of labels from the given vertex to the colored vertex above it, and is therefore an algebraic torus. The torus $T(\Gamma)$ acts on $X(\Gamma)$ by multiplication with a dense orbit.

Example 3.2. The tree Γ in Figure 1 is a colored tree with $n = 4, g = 3, m = 2$. The space of balanced labellings is

$$X(\Gamma) = \{(x_1, ..., x_6) \in \mathbb{C}^6 \mid x_1x_3 = x_1x_4 = x_2x_5 = x_2x_6, x_3 = x_4, x_5 = x_6\}$$

and admits an action of the torus $T(\Gamma) = \{(x_1, ..., x_6) \in X(\Gamma) \mid x_i \neq 0\} \simeq (\mathbb{C}^*)^3$.

Proposition 3.3. [9] *There exists an isomorphism of a Zariski open neighborhood of $M_{n,1,\Gamma}$ in $M_{n,1,\Gamma} \times X(\Gamma)$ with a Zariski open neighborhood of $M_{n,1,\Gamma}$ in $\overline{M}_{n,1}(\mathbb{A})$.*



FIGURE 1. A colored tree

Recall that affine toric varieties are classified by finitely generated cones [2]. We wish to give a description of the cone $C(\Gamma)$ that induces the toric variety $X(\Gamma)$. Note that the part of Γ below the colored vertices does not affect $X(\Gamma)$. Hence, for the rest of this section, it suffices to assume that the colored tree Γ that does not contain any vertex below any colored vertex. Recall from e.g. [2, p. 18] that the coordinate ring of $X(\Gamma)$ is generated by symbols f_μ for $\mu \in C(\Gamma)^\vee$, modulo the ideal $I(\Gamma)$ generated by relations

$$f_{\mu_1}^{\alpha_1}...f_{\mu_N}^{\alpha_N} - f_{\mu_1}^{\beta_1}...f_{\mu_N}^{\beta_N} \tag{10}$$

where $\alpha_1, ..., \alpha_N, \beta_1, ..., \beta_N$ are non negative integers satisfying

$$\alpha_1\mu_1 + ... + \alpha_N\mu_N = \beta_1\mu_1 + ... + \beta_N\mu_N.$$

We define a labelling of the edges by weights recursively as follows. First consider the case that Γ is a tree with one non-colored vertex. Label the edges below the vertex v_i by some variable e_i^*. Next suppose Γ has g non-colored vertices. By induction, assume that we have labelled the edges of the *principal subtrees* $\Gamma_1, ..., \Gamma_m$ of Γ, that is, the trees $\Gamma_1, ..., \Gamma_m$ which correspond to the vertices adjacent to the vertex in the root edge. We have thus labelled all the edges of Γ except for the principal branches. Denote by $s(\Gamma)$ the sum of the labels of the edges of a non-self-crossing path from the principal vertex to a colored vertex. Denote $s_k = s(\Gamma_k)$ for each tree Γ_k. If v_g is the principal vertex, let

$$s = s(\Gamma) = e_g^* + s_1 + ... + s_m. \tag{11}$$

Label the principal branch d_i with

$$w_{d_i} = s - s_i = e_g^* + \sum_{j \neq i} s_j. \tag{12}$$

All the edges d of Γ are now labelled by weights w_d.

Example 3.4. Figure 2 illustrates the labels of the edges of Γ.

Using (12), one sees that $s = s(\Gamma)$ is the sum of the labels of the edges of a non-self-crossing path from the principal vertex v_0 to a colored vertex and s is independent of the path chosen. Define $\chi(\Gamma)$ to be the convex cone generated by the weights w_d,

$$\chi(\Gamma) = \text{hull } \{f^{w_d} \mid d \in \mathrm{E}(\Gamma)\} = \text{hull } \bigcup_{j=1}^{m} \chi(\Gamma_j) \cup \{f^{w_{d_i}} \mid 1 \leq i \leq m\}. \tag{13}$$



Figure 2. An example of a labelling

Theorem 3.5. *There is a canonical isomorphism from* $X(\Gamma)$ *to* $\mathrm{Spec}(\mathbb{C}[\chi(\Gamma)])$.

The proof will be given after the following lemma.

Definition 3.6. Suppose A, B are two disjoint subsets of $\mathrm{E}(\Gamma)$. We write $A \sim B$ if there exists a vertex and two non-self-crossing paths γ_1 and γ_2 from that vertex to some two colored vertices so that A and B respectively contain exactly the edges of the paths γ_1 and γ_2.

Example 3.7. The set $A=\{x_1,x_3\}$ is equivalent to $B=\{x_2,x_6\}$ in example above.

Lemma 3.8. *Suppose* A *and* B *are two disjoint multisets of elements of* $\mathrm{E}(\Gamma)$. *Then*

$$\sum_{d\in A} w_d = \sum_{d'\in B} w_{d'} \tag{14}$$

if and only if A *and* B *can be partitioned into disjoint unions of* $\{A_1,...,A_r\}$ *and* $\{B_1,...,B_r\}$ *where* $A_l \sim B_l$ *for* $1\le l\le r$.

Example 3.9. In the example above, $A=\{3x_1,2x_3,x_4,x_5\}$ and $B=\{3x_2,4x_6\}$ satisfy (14). Hence, we can write $A_1=\{x_1,x_3\}\sim B_1=\{x_2,x_6\}, A_2=\{x_1,x_3\}\sim B_2=\{x_2,x_6\}, A_3=\{x_1,x_4\}\sim B_3=\{x_2,x_6\}, A_4=\{x_5\}\sim B_4=\{x_6\}$.

Proof. The only if part is already discussed. We only need to show the other direction. As before, it suffices to consider the case that there are no non-colored vertices below the colored vertices. When the number of non-colored vertices is 1 , the statement of the lemma is trivial. Assume the proposition holds for any tree with number of vertices less than g. Consider a tree Γ with g non-colored vertices. Denote by $\alpha_1,...,\alpha_m$ and $\beta_1,...,\beta_m$ the multiplicities of the principal branches $d_1,...,d_m$ in A and B. Since $A\cap B=\emptyset$, we have $\alpha_i\beta_i=0$. Equation (14) implies

$$\sum_{i=1}^m \alpha_i = \sum_{i=1}^m \beta_i, \quad -\alpha_i s_i + \sum_{d\in A\cap\Gamma_i} w_d = -\beta_i s_i + \sum_{d'\in B\cap\Gamma_i} w_{d'} \tag{15}$$

for every $1\le i\le m$. Without loss of generality, we can assume $\beta_i=0$. Then

$$\sum_{d\in A\cap\Gamma_i} w_d = \alpha_i s_i + \sum_{d'\in B\cap\Gamma_i} w_{d'}.$$

By induction, we may assume that there exists a partition $A \cap \mathrm{E}(\Gamma_i)$ and $B \cap \mathrm{E}(\Gamma_i)$ into $\{A^i_1, ..., A^i_{r_i}\}$ and $\{B^i_{\alpha_i+1}, ..., B^i_{r_i}\}$ such that for $1 \leq j \leq \alpha_i$, A^i_j contains exactly all edges of a non-self-crossing path from v_i to a colored vertex. Moreover, for $\alpha_i + 1 \leq j \leq r_i$,

$$A^i_j \sim B^i_j. \tag{16}$$

Since A contains α_i principal branch d_i, we can add one edge d_i in each A^i_j for every $1 \leq j \leq \alpha_i$. After this modification, $A^i_j \sim B^k_l$ for $1 \leq j \leq \alpha_i + \beta_i$ and $1 \leq l \leq \alpha_k + \beta_k$. Then (15) and (16) imply that we can partition A and B into $\{A_1, ..., A_r\}$ and $\{B_1, ..., B_r\}$ such that $A_i \sim B_i$ for every $1 \leq i \leq r$. □

Proof of Theorem 3.5. Let $A = \{\alpha_1 w_{d_1}, ..., \alpha_N w_{d_N}\}$ and $B = \{\beta_1 w_{d_1}, ..., \beta_N w_{d_N}\}$. Lemma 3.8 yields that A and B can be partitioned into $A_1, ..., A_r, B_1, ..., B_r$ so that $A_i \sim B_i$ for $1 \leq i \leq r$. Thus, by the description (10) of the relations, $\mathrm{Spec}(\mathbb{C}[\chi(\Gamma)]) = X(\Gamma)$. □

It follows from the theorem that the cone $C(\Gamma)$ corresponding to the toric variety $X(\Gamma)$ is the cone dual to the rational span of $\chi(\Gamma)$. Next we find a minimal set $G(\Gamma)$ of generators of $C(\Gamma)$ by an inductive argument on the number of vertices $g(\Gamma)$ of Γ.

(a) If $g(\Gamma) = 1$, then $G(\Gamma) = e_1$.
(b) If $g > 1$, then $G(\Gamma) = \{e_g + \alpha_1(w_1 - e_g) + ... + \alpha_m(w_m - e_g) \mid w_i \in G(\Gamma_i), \alpha_i \in \{0, 1\}\}$.

The cone $C(\Gamma)$ is g-dimensional and the vectors in $G(\Gamma)$ are in $\mathbb{Z}^g \subset \mathbb{R}^g$.

Theorem 3.10. *$G(\Gamma)$ is a minimal set of generators of $C(\Gamma) = (\mathrm{hull}_{\mathbb{Q}}\, \chi(\Gamma))^\vee$.*

Example 3.11. The tree Γ in Figure 1 can be split into two principal subtrees Γ_1 and Γ_2, as illustrated in Figure 2. Since $G(\Gamma_1) = \{e_1\}$ and $G(\Gamma_2) = \{e_2\}$, we obtain $G(\Gamma) = \{e_1, e_2, e_3, e_1 + e_2 - e_3\}$. The cone generated by $\{e_1, e_2, e_3, e_1 + e_2 - e_3\}$ is the cone $C(\Gamma)$ corresponding to the toric variety $X(\Gamma)$.

Proof of Theorem 3.10. We must show that $C(\Gamma) = \tilde{C}(\Gamma)$, where $\tilde{C}(\Gamma)$ is the cone generated by vectors in $G(\Gamma)$. First note that with s as in (11), for every $v \in G(\Gamma)$,

$$\langle s, v \rangle = 1. \tag{17}$$

This follows by induction on the number of vertices from the observation that

$$\langle s, v \rangle = 1 + \sum_{i=1}^{m} \alpha_i(\langle s_i, v_i \rangle - 1).$$

Next we show $C(\Gamma) \subset \tilde{C}(\Gamma)$. For $v \in G(\Gamma)$ and $w \in \chi(\Gamma)$, either $\langle w, v \rangle = \alpha_i \langle w, v_i \rangle$ if $w \in G(\Gamma_i)$ or $\langle w, v \rangle = \langle s - s_i, v \rangle = \langle s, v_i \rangle - \alpha_i \langle s_i, v_i \rangle = 1 - \alpha_i$ if $w = w_{d_i}$ for some $1 \leq i \leq m$. Hence $\langle w, v \rangle \geq 0$ and $C(\Gamma) \subset \tilde{C}(\Gamma)$.

Given $v \in \tilde{C}(\Gamma)$, we now show that v is a non-negative linear combination of elements of $G(\Gamma)$. For $g = 1$, the claim is trivial. Assume the claim is true for all trees with less than g vertices. Let Γ be a tree with g vertices. Since $\chi(\Gamma_i) \subset \chi(\Gamma)$, we can write v

uniquely $v = -\beta e_g + \sum_{i=1}^{m} z_i$ where $\beta \in \mathbb{R}$ and $z_i \in C(\Gamma_i) = \tilde{C}(\Gamma_i)$. Write $z_i = \sum_{v \in G(\Gamma_i)} \lambda_v^{(i)} v$ where $\lambda_v^{(i)} \geq 0$. Hence we can write

$$v = -\beta e_g + \sum_{i=1}^{m} \sum_{v \in G(\Gamma_i)} \lambda_v^{(i)} v.$$

If $\beta \leq 0$ then we are done since $e_g \in G(\Gamma)$. If $\beta > 0$, denote by

$$\lambda_i = \sum_{v \in G(\Gamma_i)} \lambda_v^{(i)} \geq 0.$$

Then by (17), $\langle w_{d_i}, v \rangle = -\beta + \sum_{j \neq i} \lambda_j \geq 0$.

To write v as a non-negative linear combination of elements in $\chi(\Gamma)$, we proceed as follows. Without loss of generality, assume $\lambda_m = \min\{\lambda_1, ..., \lambda_m\}$. If $\lambda_m = 0$, then move to the second minimum. If $\lambda_m > 0$, for $1 \leq i \leq m-1$, split each sum

$$z_i = \sum_{v \in G(\Gamma_i)} \gamma_v^{(i)} + \sum_{v \in G(\Gamma_i)} \delta_v^{(i)} v$$

where $\gamma_v^{(i)}, \delta_v^{(i)} \geq 0, \gamma_v^{(i)} + \delta_v^{(i)} = \lambda_v^{(i)}$ and $\sum_{v \in C(\Gamma_i)} \delta_v^{(i)} = \lambda_m$. Since

$$\sum_{v \in C(\Gamma_i)} \delta_v^{(i)} = \sum_{v \in C(\Gamma_m)} \lambda_v^{(m)} = \lambda_m,$$

we can write

$$-(m-1)\lambda_m e_g + \sum_{i=1}^{m-1} \big(\sum_{v \in G(\Gamma_i)} \delta_v^{(i)} v \big) + \sum_{v \in G(\Gamma_m)} \lambda_v^{(m)} v$$

as non-negative linear combination of elements of $G(\Gamma)$. If $-\beta' = -\beta + (m-1)\lambda_m \geq 0$, we are done. Otherwise, reapply the process to

$$-\beta' e_g + \sum_{i=1}^{m-1} \sum_{v \in G(\Gamma_i)} \gamma_v^{(i)} v.$$

This is justified since

$$-\beta' + \sum_{j \neq i, j=1}^{m-1} \gamma_j = -\beta + (m-1)\lambda_m + \sum_{j \neq i, j=1}^{m-1} (\lambda_j - \lambda_m) = -\beta + \sum_{i=1}^{m-1} \lambda_i \geq 0.$$

This process has to stop at some time and therefore we obtain v has to be a non-negative linear combination of elements in $G(\Gamma)$. Hence $C(\Gamma) = \tilde{C}(\Gamma)$.

It remains to show that $G(\Gamma)$ is a minimal set of generators of $C(\Gamma)$. We argue by induction: suppose that $G(\Gamma_i)$ is a minimal set of generators for each $C(\Gamma_i)$. Let $v \in G(\Gamma)$ be non-negative linear combination of other elements in $G(\Gamma)$. Then projecting onto the space spanned by $G(\Gamma_i)$ gives a violation of the inductive hypothesis. $\square$

The following corollaries follow immediately from Theorem 3.10:

Corollary 3.12. *The dimension of* $X(\Gamma)$ *equals the number of non-colored vertices* $g(\Gamma)$.

Corollary 3.13. *If the number of 1 dimensional faces of* $C(\Gamma_i)$ *is* r_i *for* $1 \leq i \leq m$, *then the number of 1 dimensional faces of* $C(\Gamma)$ *is* $r = (r_1 + 1)...(r_m + 1)$.

4. LOCAL DESCRIPTION OF CARTIER BOUNDARY DIVISORS

Recall the description of invariant Weil divisors of an affine toric variety $X(\Gamma)$ with cone $C(\Gamma)$ [2]:

Proposition 4.1. *There is a bijection between the one dimensional faces of* $C(\Gamma)$ *with the invariant prime Weil divisors of* $X(\Gamma)$.

We first identify the invariant prime Weil divisors of $X(\Gamma)$.

Definition 4.2. A subset $Y = \{y_1, ..., y_r\} \subset \mathrm{E}(\Gamma)$ is called *complete* if it has the following property: when we set $x_{y_1} = \cdots = x_{y_r} = 0$, the relation $x_k...x_l = 0$ holds for some edges $k, ..., l \in \mathrm{E}\,\Gamma$ if and only if at least one of the edges in the product is in Y.

Definition 4.3. A complete subset Y of $\mathrm{E}(\Gamma)$ is called *minimally complete* if it does not properly contain any complete subset of $\mathrm{E}(\Gamma)$.

Denote by $\mathcal{D}(\Gamma)$ the set of minimally complete subsets $Y \subset \mathrm{E}(\Gamma)$. A minimally complete subset can be characterized combinatorially as follows.

Example 4.4. The minimally complete subsets of $\mathrm{E}(\Gamma)$, where Γ is the tree in Figure 1, are $\{x_1, x_2\}$, $\{x_1, x_5, x_6\}$, $\{x_2, x_3, x_4\}$, $\{x_3, x_4, x_5, x_6\}$.

Lemma 4.5. *A subset* $Y \subset \mathrm{E}(\Gamma)$ *is minimally complete if and only if each non-self crossing path from* v_0 *to a colored vertex contains exactly one edge in* Y.

Proof. Suppose Y is minimally complete and $y \in Y$. If v_k is the upper endpoint of y, then all the variables of the edges above v_k are not in Y. An induction argument implies any path from v_0 to a colored vertex contains exactly one edge in Y. The proof of the converse is similar. □

Corollary 4.6. *If the number of minimally complete subsets of* $\mathrm{E}(\Gamma_i)$ *is* r_i, *then the number of minimally complete sets of* $\mathrm{E}(\Gamma)$ *is* $r = (r_1 + 1)...(r_m + 1)$.

From Corollary 3.13, we also obtain

Corollary 4.7. *The number of 1 dimensional faces of* $C(\Gamma)$ *equals the number of minimally complete subsets of* $\mathrm{E}(\Gamma)$.

We can now describe the set of invariant Weil divisors of $X(\Gamma)$ as follows.

Proposition 4.8. *There is a bijection between the set of invariant prime Weil divisors and the set of minimally complete subsets of* $\mathrm{E}(\Gamma)$. *More explicitly, each invariant Weil divisor has the form* $D_Y = \{(x_1, ..., x_N) \in X(\Gamma) \mid x_i = 0\, \forall x_i \in Y\}$ *for some minimally complete subset* Y.

Proof. Suppose D is an invariant Weil divisor of $X(\Gamma)$. Then D is the union of some orbits. Given a point $(x_1, ..., x_N) \in D$, define

$$Y = \{i \mid x_i = 0, 1 \leq i \leq N\}.$$

The orbit of this point under the torus action is

$$\tilde{D}_Y = \{(x_1, ..., x_N) \in X(\Gamma) \mid x_i = 0\, \forall i \in Y \;\text{ and }\; x_k \neq 0\, \forall k \notin Y\}.$$

Since D is a subvariety, D must contain the Zariski closure of $\tilde{D}_Y$, which is D_Y.

We now show if D_Y is a subvariety of codimension 1, then Y is minimally complete. We notice that given a minimal complete subset $Y \subset \mathrm{E}(\Gamma)$, for each principal subtree Γ_i, either $x_{d_i} = 0$ or D_Y induces a minimally complete subset Y_i of $\mathrm{E}(\Gamma_i)$. Hence the dimension of D_Y is $g(\Gamma_1) + ... + g(\Gamma_m) = g - 1$. Thus D_Y is a subvariety of $X(\Gamma)$ of codimension 1. From this argument, it also follows immediately that if Y isn't minimally complete, then D_Y is a subvariety of $X(\Gamma)$ of codimension at least 2. Hence, the torus invariant subvarieties of $X(\Gamma)$ codimension 1 are exactly all D_Y, where $Y \subset \mathrm{E}(\Gamma)$ is minimally complete.

It is left to show that each D_Y is prime. From Proposition 4.1 and Proposition 4.7, the number of Weil divisors equals the number of 1 dimensional faces of $C(\Gamma)$ which equals the number of minimally complete subset of $\mathrm{E}(\Gamma)$. Therefore the invariant Weil divisors of $X(\Gamma)$ are exactly all D_Y, where $Y \subset \mathrm{E}(\Gamma)$ is minimally complete. □

Example 4.9. The invariant prime Weil divisors of $X(\Gamma)$ from Figure 1 are $D_{\{1,2\}}, D_{\{1,5,6\}}, D_{\{2,3,4\}}, D_{\{3,4,5,6\}}$.

We describe the rays of $C(\Gamma)$ corresponding to minimally complete subsets. Let $D = D_Y$ be Weil divisor as above. Unless $x_{d_i} \in Y$, the principal subtree Γ_i has an induced Weil divisor $D_{Y_i} \subset V(\Gamma_i)$.

Proposition 4.10. *Let* D_Y, D_{Y_i} *be as above. Assume that the corresponding 1 dimensional face of* D_{Y_i} *is generated by* $v_i \in G(\Gamma_i) \subset G(\Gamma)$. *Let* $I = \{i \mid x_{d_i} \notin Y, 1 \leq i \leq m\}$. *The corresponding 1 dimensional face* τ *of* $C(\Gamma)$ *is generated by*

$$v_Y = e_g + \sum_{i \in I}(v_i - e_g). \tag{18}$$

Proof. Denote $\tau_{\mathbb{Z}}^{\perp} = \tau^{\perp} \cup \mathrm{Hom}(\mathbb{Z}^g, \mathbb{Z})$ and $\tau_{\mathbb{Z}}^{\vee} = \tau^{\vee} \cup \mathrm{Hom}(\mathbb{Z}^g, \mathbb{Z})$. Let $\phi \in \mathrm{Hom}(\tau_{\mathbb{Z}}^{\perp} \cap M, \mathbb{C}^*)$ and $\tilde{\phi} \in \mathrm{Hom}(\tau_{\mathbb{Z}}^{\vee} \cap M, \mathbb{C})$ be the image of ϕ in the zero extension map $\mathrm{Hom}(\tau_{\mathbb{Z}}^{\perp} \cap M, \mathbb{C}^*) \to \mathrm{Hom}(\tau_{\mathbb{Z}}^{\vee} \cap M, \mathbb{C})$. Notice $i \in I$ if and only if $\tilde{\phi}(v_i) = 0$. By induction, $\tilde{\phi}$ corresponds to the point $(a_1, ..., a_N) \in X(\Gamma)$ where $a_j = 0$ if and only if either $x_j \in Y_i$

for some $i \in I$ or $x_j = x_{d_i}$ for some $i \notin I$. Hence the closure of the orbit $O(\tau)$ is exactly D_Y. □

We recall the classification of Cartier divisors of a toric variety X with cone C.

Theorem 4.11. *The divisor* $\sum\limits_{Y \in \mathcal{D}(\Gamma)} a_Y D_Y$ *is a Cartier divisor if and only if there exists* $u \in M = \mathrm{Hom}(\mathbb{Z}^n, \mathbb{Z})$ *such that* $\langle u, v_Y \rangle = a_Y$. [2]

For each vertex v_k of Γ, we define

$$\mathcal{D}_k = \{D_Y \mid Y \text{ contains some edges of } \Gamma_k\}, \quad D_k = \sum_{D \in \mathcal{D}_k} D.$$

Hence if $D_Y \in \mathcal{D}_k$, Y does not contain edges of Γ which are above v_k.

Proposition 4.12. *The group of invariant Cartier divisors is generated by* $D_0, ..., D_{g-1}$.

Example 4.13. The set of Cartier divisors of $X(\Gamma)$, where Γ is the tree in Figure 1, is generated by

$$\{D_{\{1,2\}} + D_{\{1,5,6\}} + D_{\{2,3,4\}} + D_{\{3,4,5,6\}}, D_{\{2,3,4\}} + D_{\{3,4,5,6\}}, D_{\{1,5,6\}} + D_{\{3,4,5,6\}}\}.$$

Thus $n_{\{1,2\}}D_{\{1,2\}} + n_{\{3,4,5,6\}}D_{\{3,4,5,6\}} + n_{\{1,5,6\}}D_{\{1,5,6\}} + n_{\{2,3,4\}}D_{\{2,3,4\}}$ is a Cartier divisor if and only if $n_{\{1,2\}} + n_{\{3,4,5,6\}} = n_{\{1,5,6\}} + n_{\{2,3,4\}}$.

Proof of Proposition 4.12. We first check that D_k is a Cartier divisor. Recall the notation in (11), $s_k = s(\Gamma_k) \in \mathrm{Hom}(\mathbb{Z}^g, \mathbb{Z})$ for each vertex v_k. Note that s_k satisfies $\langle s_k, v_Y \rangle = 1$ if $Y \in \mathcal{D}_k$ and $\langle s_k, v_Y \rangle = 0$ otherwise. Indeed, for each $Y \in \mathcal{D}_k$, D_Y induces another Cartier divisor D_{Y_k} and by (17), we have $\langle s_k, v_{Y_k} \rangle = 1$. This implies $\langle s_k, v_Y \rangle = 1$. On the other hand, if $Y \notin \mathcal{D}_k$, then Y does not contain any edges in Γ_k. Thus, $\langle s_k, v_Y \rangle = 0$. Hence D_k is a Cartier divisor. Moreover, $s_k - \sum\limits_{v_k v_i \in \mathrm{B}(\Gamma_k)} s_i = {e_k}^*$. Hence, $\{s_k\}_{0 \le k \le g-1}$ generates M. Therefore, $D_0, ..., D_{g-1}$ generates the set of Cartier divisors of $X(\Gamma)$. □

We have the following description of the subspace of Cartier divisors of $X(\Gamma)$.

Theorem 4.14. $\sum\limits_D n_D D$ *is a Cartier divisor of* $X(\Gamma)$ *if and only if* $\sum\limits_D m_D n_D = 0$ *for every* $(m_D)_D \in \mathbb{Z}^W$ *that satisfies* $\sum\limits_{Y : i \in Y} m_{D_Y} = 0$ *for every edge* $i \in \mathrm{E}(\Gamma)$. *Here* W *is the number of prime Weil invariant divisors of* $X(\Gamma)$.

Proof. The set of integral Cartier divisors of $X(\Gamma)$ forms a sublattice N of the space of integral Weil divisors $\mathbb{Z}^W$. Since $D_0, ..., D_{g-1}$ are $\mathbb{R}$- linearly independent, $N_{\mathbb{R}}$ has dimension g. Suppose $(m_D)_D \in \mathbb{Z}^W$ such that $\sum\limits_{Y : i \in Y} m_{D_Y} = 0$ for all i. Consider a non-self-crossing path from a vertex v_k to a colored vertex. By summing over edges of the path, we obtain $\sum\limits_{D \in \mathcal{D}_k} m_D = \sum\limits_{i=1}^{j} \sum\limits_{Y : i \in Y} m_{D_Y} = 0$. Thus, if $\sum\limits_D n_D D$ is a Cartier divisor, then

D is a combination of $D_0, \ldots, D_{g-1}$ and so $\sum_D m_D n_D = 0$. On the other hand, the set of $(m_D)_D \in \mathbb{Z}^W$ that satisfies the condition in the Theorem forms a sublattice $\tilde{N}$ with dimension at least $W - g$, since the conditions are linearly independent. Therefore, the dimension of $N_{\mathbb{R}}$, which is a subspace of $\tilde{N}_{\mathbb{R}}^{\perp}$, is at most g. Combining these observations shows $N_{\mathbb{R}} = \tilde{N}_{\mathbb{R}}^{\perp}$.

Suppose that $(n_D)_D \in \mathbb{Z}^W$ satisfies the condition in the Theorem. By the previous paragraph,

$$\sum_D n_D D = \frac{1}{s} \sum_{i=0}^{g-1} r_i D_i$$

for some integers r_i and s such that $s > 0$. It is suffices now to show that $s | r_i$ for every $0 \leq i \leq g-1$. To see this, note that

$$\sum_D n_D D = \frac{1}{s} \left(\sum_{i=0}^{g} (r_i \sum_{D \in \mathcal{D}_i} D) \right) = \frac{1}{s} \left(\sum_D (\sum_{i: D \in \mathcal{D}_i} r_i) D \right).$$

Thus

$$n_D = \frac{1}{s} \sum_{i: D \in \mathcal{D}_i} r_i.$$

For the principal vertex v_0, define $D^0 = D_Y$ where $Y = \{d_1, ..., d_m\}$. More generally, for any vertex v_k, let γ_k be the down-path from v_0 to v_k, and let D^k to be the divisor $D^k = D_Y$ where Y is the set of edges immediately below the vertices in γ_k. Then

$$n_{D^k} = \frac{1}{s} \sum_{v_i \in \gamma_k} r_i \in \mathbb{Z} \tag{19}$$

Since $n_{D^0} \in \mathbb{Z}$, we obtain $s | r_0$. Induction on the length of the path γ_k gives $s | r_k$. ☐

5. Global description of Cartier boundary divisors

In this section we give an explicit criterion for a divisor D of the form (5) to be Cartier. It is easy to see from the local description of the moduli space given below that any divisor of type I is Cartier, so it suffices to consider divisors of type II. To describe the answer, let $I = \{1, \ldots, n\}$, $\mathrm{Par}(I)$ the set of *non-trivial* partitions of I, and $\mathcal{P}(I)$ the power set of *non-empty* proper subsets of I. We identify the set of prime Weil boundary divisors of type I with the subset of elements of $\mathcal{P}(I)$ of size at least two, and the prime Weil boundary divisors of type II with $\mathrm{Par}(I)$. Thus in particular the space of Weil boundary divisors becomes identified with $\mathbb{Z}[\mathrm{Par}(I)]$, by the map

$$\left\{ \sum_P n(P) D_P \right\} \to \mathbb{Z}[\mathrm{Par}(I)], \quad \sum_P n(P) D_P \mapsto n.$$

Let $Z(I)$ denote the natural incidence relation,

$$Z(I) = \{(S, P) \in \mathcal{P}(I) \times \mathrm{Par}(I) \,|\, S \in P\}.$$

Let $p : Z(I) \to \mathcal{P}(I), \quad q : Z(I) \to \mathrm{Par}(I)$ denote the projections. We have a natural map from the space of functions on $\mathrm{Par}(I)$ to functions on $\mathcal{P}(I)$ given by pullback and push-forward:

$$p_* q^* : \mathbb{Z}[\mathrm{Par}(I)] \to \mathbb{Z}[\mathcal{P}(I)], \quad (p_* q^* h)(S) = \sum_{S \in P} h(P).$$

A *relations* on the space of Cartier divisors is a collections of coefficients $\{m_{I_1,\dots,I_r}\} \in \mathbb{Z}[\mathrm{Par}(\{1,\dots,n\})]$ such that

$$\sum_{I_1,\dots,I_r} m_{I_1,\dots,I_r} n_{I_1,\dots,I_r} = 0$$

for every Cartier divisor $D = \sum_{I_1,\dots,I_r} n_{I_1,\dots,I_r} D_{I_1,\dots,I_r}$. The space of relations forms a subgroup of $\mathbb{Z}[\mathrm{Par}(I)]$. The main result of this note is the following:

Theorem 5.1. *The group of relations on the space of Cartier divisors of type II is the kernel of $p_* q^*$. The space of Cartier divisors of type II is the image of $\mathbb{Z}[\mathcal{P}(I)]$ under $q_* p^*$.*

Example 5.2. For $n = 2$ there are two boundary divisors and no relations. For $n = 3$ there are six boundary divisors and no relations. For $n = 4$ there are $|\mathcal{P}(\{1,2,3,4\})| - 4 = 11$ boundary divisors of type I, and $|\operatorname{Par}(\{1,2,3,4\})| = 1+6+3+4 = 14$ boundary divisors of type II. A divisor

$$D = \sum n_{I_1,\dots,I_r} D_{I_1,\dots,I_r}$$

is Cartier only if the three relations (as i,j,k,l vary)

$$n_{\{i,j\},\{k\},\{l\}} + n_{\{i\},\{j\},\{k,l\}} - n_{\{i,j\},\{k,l\}} - n_{\{i\},\{j\},\{k\},\{l\}} \tag{20}$$

hold. Thus the space of Cartier boundary divisors of type II is a 11-dimensional subspace of the space of the 14-dimension space of Weil boundary divisors of type II.

Let Γ be a colored tree, and choose a bijection from the set of colored vertices of Γ to $I = \{1, ..., n\}$.

Definition 5.3. A tree Γ is *simple* if it has a single vertex.

For each partition $\{I_1, ..., I_r\}$ of $I = \{1, ..., n\}$, define the tree $\Gamma_{I_1,\dots,I_r}$ as follows: $\Gamma_{I_1,\dots,I_r}$ has r principal subtrees which are respectively the simple trees whose colored vertices are those in I_r. Given $v, \tilde{v} \in \mathrm{Vert}(\Gamma)$, we write $vE\tilde{v}$ if there is an edge connecting v and $\tilde{v}$.

Definition 5.4. A *tree homomorphism* $f : \mathrm{Vert}(\Gamma) \to \mathrm{Vert}(\Gamma')$ is a map that maps the vertices and edges of Γ to the vertices and edges of Γ' respectively and satisfies:

(a) f maps the principal vertex v_0 of Γ to the principal vertex v_0' of Γ'.
(b) If $v, \tilde{v} \in \mathrm{Vert}(\Gamma)$ satisfies $vE\tilde{v}$, then $f(v)Ef(\tilde{v})$.
(c) f maps bijectively the set of colored vertices of Γ to the set of colored vertices of Γ'.

Definition 5.5. A partition $\{I_1, ..., I_r\}$ of $\{1, ..., n\}$ is *compatible* with Γ if there exists a tree homomorphism $f : \Gamma \to \Gamma_{I_1,...,I_r}$. The set of compatible partitions of Γ is denoted by $\mathrm{Par}(\Gamma)$.

Denote by $\mathrm{Par}(\Gamma)$ the set of compatible partitions of Γ.

Proposition 5.6. *There is a bijection between the set of compatible partitions and the set of minimally complete subsets of* $\mathrm{E}(\Gamma)$*.*

Example 5.7. For the tree Γ in Figure 1, the correspondence between the minimally complete subsets of $\mathrm{E}(\Gamma)$ and the compatible partitions of $\{1, ..., n\}$ is

$$\{x_1, x_2\} \longleftrightarrow \{1, 2\}, \{3, 4\}, \quad \{x_1, x_5, x_6\} \longleftrightarrow \{1, 2\}, \{3\}, \{4\}$$
$$\{x_2, x_3, x_4\} \longleftrightarrow \{1\}, \{2\}, \{3, 4\}, \quad \{x_3, x_4, x_5, x_6\} \longleftrightarrow \{1\}, \{2\}, \{3\}, \{4\}.$$

Proof. When $g = 1$, there is one minimally complete subset which corresponds to the partition $\{\{1\}, ..., \{n\}\}$. Given a minimally complete subset $Y = \{y_1, ..., y_r\}$. By cutting the edge y_k, we split the tree into two parts. Denote by I_k the set of colored points in the part not containing the root of Γ. Then $I_1, ..., I_r$ is a nontrivial partition of $\{1, ..., n\}$. We can construct a homomorphism $f : \Gamma \to \Gamma_{I_1,...,I_r}$ as follows. For each principal subtree Γ_j of Γ, the partition $\{I_1, ..., I_r\}$ induces a partition P_j of the set of colored points of Γ_j. In fact, the induced partition is a subset of the partition $\{I_1, ..., I_r\}$. Hence there is a natural inclusion map $i_j : \Gamma_{P_j} \to \Gamma_{I_1,...,I_r}$. If P_j is trivial, define $f_j : \Gamma_j \to \Gamma_{P_j}$ to be the identity map. Otherwise, by induction hypothesis, we obtain a homomorphism $f_j : \Gamma_j \to \Gamma_{P_j}$ for each principal subtree Γ_j. The map $f : \Gamma \to \Gamma_{I_1,...,I_r}$ such that $f = i_j \circ f_j$ when restricted to each principal subtree Γ_j is a tree homomorphism we need.

Let $\{I_1, ..., I_r\}$ be a compatible partition with a tree homomorphism $f : \Gamma \to \Gamma_{I_1,...,I_r}$. We can construct the corresponding minimally complete subset Y as follows: for each partition I_k, define y_k to be first common edge of the paths from the colored vertices in I_k to the root. Let $Y = \{y_1, .., y_r\}$. Note that Y is a minimally complete subset of the set of edges of Γ. Indeed, suppose there exist two edges y_k and y_j in Y and y_j is below y_k. Then $I_j \cap I_k \neq \emptyset$ because of the condition f maps the principal vertex v_0 of Γ to the principal vertex v_0' of $\Gamma_{I_1,..,I_r}$. It is not hard to verify that the constructions above are inverse to each other. Therefore,we obtain a bijective correspondence between the set of compatible partitions and the minimally complete subsets of $\mathrm{E}(\Gamma)$. □

From Proposition 5.6 we obtain a bijective correspondence between compatible partitions and the Weil divisors of $X(\Gamma)$. For each compatible partition $\{I_1, ..., I_r\}$ of Γ, denote by $D_{I_1,...,I_r}$ the corresponding invariant prime Weil divisors of $X(\Gamma)$. We prove the main result:

Proof of Theorem 5.1. Given $S \subset \{1, ..., n\}$. From Theorem 4.14, we obtain

$$\sum_{\{I_1,...,I_r\} \in \mathrm{Par}(\Gamma)} n_{I_1,...,I_r} D_{I_1,...,I_r}$$

is a Cartier divisor of $X(\Gamma)$ if and only if $(n_{I_1,...,I_r})_{\{I_1,...,I_r\}\in \mathrm{Par}(\Gamma)}$ is in the orthogonal complement of the kernel of $p_*q_\Gamma^*$, where

$$p_*q_\Gamma^* : \mathbb{Z}[\mathrm{Par}(\Gamma)] \to \mathbb{Z}[\mathcal{P}(\Gamma)], \quad (p_*q_\Gamma^*(m))(S) = \sum_{S\in\{I_1,...,I_r\}} m_{I_1,...,I_r}.$$

Note that there is a natural embedding of $\mathrm{Ker}\, p_*q_\Gamma^*$ in $\mathrm{Ker}\, p_*q^*$ by the map $\pi_\Gamma : \mathrm{Ker}\, p_*q_\Gamma^* \to \mathrm{Ker}\, p_*q^*$ which preserves $m_{I_1,...,I_r}$ if $\{I_1, ..., I_r\} \in \mathrm{Par}(\Gamma)$ and maps the other $m_{I_1,...,I_r}$,where $\{I_1, ..., I_r\} \notin \mathrm{Par}(\Gamma)$, to 0. The image of π_Γ is a subspace of $\mathrm{Ker}\, p_*q^*$ and

$$\sum_{I_1,...,I_r} n_{I_1,....,I_r} D_{I_1,...,I_r}$$

is a Cartier divisor of $X(\Gamma)$ if and only if $(n_{I_1,....,I_r}) \in \mathrm{coker}\, \pi_\Gamma$. Hence, $\sum\limits_{I_1,...,I_r} n_{I_1,....,I_r} D_{I_1,...,I_r}$ is a Cartier divisor of all $X(\Gamma)$ if and only if $(n_{I_1,...,I_r}) \in \bigcap\limits_\Gamma \mathrm{coker}\, \pi_\Gamma$. Thus, it suffices to show that $\mathrm{Ker}\, p_*q^* \subset \mathrm{hull}_{\mathbb{Z}}\, \mathrm{image}\, \pi_\Gamma$. We proceed by induction on the number of colored vertices. Denote by Tree_n the class of colored metric trees with n colored vertices. For $n = 1$, the statement is obvious. Denote by $\mathcal{S}_j \subset \mathrm{Tree}_n$ the subclass of trees Γ such that the principal subtree of Γ containing the colored vertex j is a simple tree with at least 2 branches. Also, denote by $\mathcal{S}_j^1 \subset \mathrm{Tree}_n$ the subclass of trees Γ such that the principal subtree containing colored vertex j is a simple tree with exactly 1 branch. Denote by $\mathrm{Par}_j(I) \subset \mathrm{Par}(I)$ the set of partitions $\{I_1, ..., I_r\}$ such that $\{j\} \notin \{I_1, ..., I_r\}$. Given $\vec{m} = (m_{I_1,...,I_r}) \in \ker(p_*q^*)$ and Γ in $\mathcal{S}_n$, denote by $\vec{m}_\Gamma$ the orthogonal projection of $\vec{m} = (m_{I_1,...,I_r})$ to the subspace $\mathrm{image}\, p_*q_\Gamma^*$. Then $\vec{v} = \vec{m} - \sum\limits_{\Gamma\in\mathcal{S}_n} \vec{m}_\Gamma$ has $\{I_1, ..., I_r\}$-coordinate equal 0 for every $\{I_1, ..., I_r\} \in \mathrm{Par}_n(I)$ and for $\{\{1, ..., n-1\}, \{n\}\}$-coordinate. For the $\{I_1, ..., I_r, \{n\}\}$-coordinate of $\vec{v}$, we abuse the notation by denoting $v_{I_1,...,I_r} = v_{I_1,...,I_r,\{n\}}$. By the inductive hypothesis, we obtain $v_{I_1,...,I_{r-1}} \in \mathrm{hull}_{\mathbb{Z},\Gamma\in\mathrm{Tree}_{n-1}}\, \mathrm{image}\, \pi_\Gamma$. On the other hand, for $\Gamma \in \mathrm{Tree}_{n-1}$, $\mathrm{image}\, \pi_\Gamma$ can be naturally embedded in $\mathrm{image}\, \pi_{\tilde{\Gamma}}$, where $\tilde{\Gamma} \in \mathcal{S}_n^1$ is obtained from Γ by attaching a principal subtree that contains the colored vertex n. Thus $\vec{v} \in \mathrm{hull}_{\mathbb{Z},\Gamma\in\mathcal{S}_n^1}\, \mathrm{image}\, \pi_\Gamma \subset \mathrm{hull}_{\mathbb{Z},\Gamma\in\mathrm{Tree}_n}\, \mathrm{image}\, \pi_\Gamma$. By induction, $\mathrm{Ker}(p_*q^*) = \mathrm{hull}_{\mathbb{Z},\Gamma\in\mathrm{Tree}_n}\, \mathrm{image}\, \pi_\Gamma$ as claimed.

Next we prove the second claim in Theorem 5.1.

Definition 5.8. A partition $P \in \mathrm{Par}(I)$ is *simple* if there exists $S \in \mathcal{P}(I)$ such that $S \in P$ and for every $i \notin S$, $\{i\} \in P$.

The number of simple partitions is $2^n - n - 1$.

Example 5.9. When $n = 4$, there are 11 simple partitions: $\{\{1\}, \{2\}, \{3\}, \{4\}\}$, $\{\{1, 2\}, \{3\}, \{4\}\}$, $\{\{1, 3\}, \{2\}\{4\}\}$, $\{\{1, 4\}, \{2\}, \{3\}\}$, $\{\{1\}\{2, 3\}\{4\}\}$; $\{\{1\}\{2, 4\}, \{3\}\}$, $\{\{1\}, \{2\}, \{3, 4\}\}$, $\{\{1, 2, 3\}\{4\}\}$, $\{\{1, 3, 4\}\{2\}\}$, $\{\{1, 2, 4\}\{3\}\}$,$\{\{1\}\{2, 3, 4\}\}$.

Consider the unique natural extension maps

$$(p_*q^*)_{\mathbb{Q}} : \mathbb{Q}[\mathrm{Par}(I)] \to \mathbb{Q}[\mathcal{P}(I)], \quad (q_*p^*)_{\mathbb{Q}} : \mathbb{Q}[\mathrm{Par}(I)] \to \mathbb{Q}[\mathcal{P}(I)].$$

From the first claim of Theorem 5.1 and the identities

$$\ker((p_*q^*)_{\mathbb{Q}}) = \operatorname{coker}((q_*p^*)_{\mathbb{Q}}), \quad \ker((q_*p^*)_{\mathbb{Q}}) = \operatorname{coker}((p_*q^*)_{\mathbb{Q}})$$

it follows that the space of rational Cartier boundary divisors is exactly the image of $(q_*p^*)_{\mathbb{Q}}$. Suppose $\sum n_P D_P$ is a Cartier divisor with integral coefficients and hence, we can find $k \in \mathbb{Q}[\mathcal{P}(I)]$ such that

$$n_P = (q_*p^*)_{\mathbb{Q}} k(P) = \sum_{S \in P} k(S).$$

Define $\tilde{k} \in \mathbb{Z}[\mathcal{P}(I)]$ as follows: pick arbitrary integral values for $\tilde{k}(\{i\})$ such that $\sum\limits_{i=1}^{n} \tilde{k}(\{i\}) = n_{\{\{1\},\ldots,\{n\}\}}$. For $S \in \mathcal{P}(I)$, let P_S denote the unique simple partition P_S such that $S \in P$. Define

$$\tilde{k}(S) = n_{P_S} - \sum_{i \notin S} \tilde{k}(\{i\}).$$

We show that $q_*p^*\tilde{k}(P) = n_P$. Let $r = \tilde{k} - k \in \mathbb{Q}[\mathcal{P}(I)]$. Then

$$p_*q^*r(P) = \sum_{S \in P}(\tilde{k}(S) - k(S)) = \sum_{S \in P}(n_{P_S} - \tilde{k}(S) - \sum_{i \notin S} \tilde{k}(\{i\}) = \sum_{S \in P}\sum_{i \notin S}(k(\{i\} - \tilde{k}(\{i\}) = 0$$

since each i appears in exactly one $S \in P$ and $\sum\limits_i (k(\{i\}) - \tilde{k}(\{i\})) = 0$. Therefore, $q_*p^*\tilde{k}(P) = q_*p^*k(P) = n_P$ and $\tilde{k} \in \mathbb{Z}[\mathcal{P}(I)]$. □

In the remainder of this section, we explore consequences of Theorem 5.1. The following is immediate:

Corollary 5.10. *The space of Cartier boundary divisors of type II in* $\overline{M}_{n,1}(\mathbb{A})$ *has rank* $(2^n - 2) - n + 1 = 2^n - n - 1$.

Note that this is the same as the rank of Cartier boundary divisors of type I of $\overline{M}_{n,1}(\mathbb{A})$. Finally we give a geometric description of the space of Cartier boundary divisors in terms of the forgetful morphism $f : \overline{M}_{n,1}(\mathbb{A}) \to \overline{M}_n$ obtained by forgetting the scaling.

Corollary 5.11. *The space of Cartier boundary divisors in* $\overline{M}_{n,1}(\mathbb{A})$ *is generated by the inverse images of Cartier divisors under* f *and the boundary divisors of type I.*

Proof. Suppose that $D_S \subset \overline{M}_n$ is the divisor corresponding to some subset $S \subset \{1, \ldots, n\}$. Using the definition of the forgetful morphism, the inverse image $f^{-1}(D_S)$ is a Cartier divisor of $\overline{M}_{n,1}(\mathbb{A})$,

$$f^{-1}(D_S) = D_S + \sum_{S \in P} D_P.$$

The second term is the image of a standard basis vector $e_S \in \mathbb{Z}[\mathbb{P}(\{1, \ldots, n\})$ under the map q_*p^*. It follows from the description above that modulo divisors of type I, these divisors generate the space of Cartier divisors of $\overline{M}_{n,1}(\mathbb{A})$, which completes the proof. □

We remark that the inverse image of the divisor $D_{\{1\},\{2\}} \subset \overline{M}_{2,1}(\mathbb{A})$ under the forgetful morphism $f_{i,j} : \overline{M}_{n,1}(\mathbb{A}) \to \overline{M}_{2,1}(\mathbb{A})$ is always Cartier.

Example 5.12. Let $n = 4$ and $i = 1, j = 4$. Then

$$\begin{aligned} f^{-1}D_{\{1\},\{2\}} = D_{\{1\},\{2\},\{3\},\{4\}} &+ D_{\{1,2\},\{3\},\{4\}} + D_{\{1,3\},\{2\},\{4\}} \\ &+ D_{\{1,2\},\{3,4\}} + D_{\{1,3\},\{2,4\}} + D_{\{1,2,3\},\{4\}} + D_{\{1\},\{2,3,4\}}. \end{aligned} \tag{21}$$

One checks easily that the three relations in (20) hold.

6. What kind of category do CohFT's form?

Naturally the definition of morphism of CohFT's above suggests the question of what sort of category CohFT's form. We will not give a complete answer to this question, but rather we will sketch a notion of *composition* of morphisms of CohFT's. This will make CohFT's into something like a category, but with additional information necessary for composition, as in Street's notion of higher categories, see for example [10]. This notion of composition plays a natural role in the *quantum reduction with stages* conjecture in [11]. The cohomological datum needed for the composition maps is associated to a moduli space of s-scaled n-marked lines as follows.

Definition 6.1. An *s-scaled, n-marked affine line* over a field k is an affine line Σ equipped with s scalings $\phi_1, \ldots, \phi_s : T\Sigma \to \Sigma \times k$ and n distinct points $z_1, \ldots, z_n$. An *isomorphism* of s-scaled, n-marked affine lines Σ_0, Σ_1 is an isomorphism $\psi : \Sigma_0 \to \Sigma_1$ preserving the scalings and markings: $\phi_i^1 \circ D\psi = (\psi \times 1) \circ \phi_i^0$ for $i = 1, \ldots, s$ and $\psi(z_j^0) = z_j^1$ for $j = 1, \ldots, n$.

Let $k = \mathbb{C}$ and let $M_{n,s}(\mathbb{A})$ denote the moduli space of s-scaled, n-marked affine lines up to automorphism. $M_{n,s}(\mathbb{A})$ has a natural compactification obtained by allowing bubbles on which a proper subset of the scalings have gone to infinity or zero.

Definition 6.2. A *s-scaled n-marked nodal curve* $(\underline{\Sigma}, \underline{z}, \underline{\phi})$ consists of

(a) A nodal curve $\underline{\Sigma} = \Sigma_0, \ldots, \Sigma_k$ where the *root component* Σ_0 is an affine line and $\Sigma_1, \ldots, \Sigma_k$ are projective lines.
(b) A collection of distinct smooth points $\underline{z} = (z_1, \ldots, z_n) \in \underline{\Sigma}$.
(c) a collection of possibly degenerate scalings $\phi_i^j : T\Sigma_i' \to \Sigma_i' \times k$, where Σ_i' is the complement of the node connecting Σ_i to the root component;

such that

(a) for any path from Σ_0 to a terminal component and for any index i, there is exactly one component $m(i)$ on which the i-th scaling is non-degenerate
(b) the components previous to $m(i)$ have zero i-th scaling and the components after that have infinite i-th scaling.
(c) For any indices i, j, the ratio between the i-th and j-th scaling is independent of the choice of component.

A s-scaled, n-marked line $\underline{\Sigma}$ is *stable* if each component with at least one non-degenerate scaling has at least one marked or nodal point, and each component with all degenerate scalings has at least two marked or nodal points. The *combinatorial type* of an s-scaled, n-marked affine line is an $s+1$-colored tree, defined as follows. Let Γ be the tree defined by $\underline{\Sigma}$. For each i, the i-colored vertices are those on which the i-th scaling is finite.

Let $\overline{M}_{n,s}(\mathbb{A})$ denote the moduli space of s-scaled, n-marked lines. The boundary of $\overline{M}_{n,s}(\mathbb{A})$ can be described as follows.

(a) For any subset $I \subset \{1, \dots n\}$ of order at least two there is a divisor

$$\iota_I : D_I \to \overline{M}_{n,s}(\mathbb{A})$$

and an isomorphism

$$\varphi_I : D_I \to \overline{M}_{0,|I|+1}(\mathbb{A}) \times \overline{M}_{n-|I|+1,s}$$

corresponding to the formation of a bubble containing the markings $z_i, i \in I$ with zero scaling on that bubble.

(b) For any partition $I_1 \cup \dots \cup I_r$ of $\{1, \dots, n\}$ of order at least two and subset $J \subset \{1, \dots, s\}$ there is a divisor

$$\iota_{I_1 \cup \dots \cup I_r, J} : D_{I_1 \cup \dots \cup I_r, J} \to \overline{M}_{n,s}(\mathbb{A})$$

with an isomorphism

$$\varphi_{I_1 \cup \dots \cup I_r, J} : D_{I_1 \cup \dots \cup I_r, J} \to \overline{M}_{r+1, s-|J|}(\mathbb{A}) \times \prod_{i=1}^{r} \overline{M}_{|I_i|+1,|J|}(\mathbb{A})$$

corresponding to the formation of r bubbles containing markings $I_j, j = 1, \dots, r$ with the scalings $j \in J$ becoming infinite on those bubbles.

The union of these divisors is the boundary of $M_{n,s}$:

$$\partial \overline{M}_{n,s} = \bigcup_{I \subset \{1,\dots n\}} D_I \cup \bigcup_{I_1 \cup \dots \cup I_r = \{1,\dots,n\}} D_{I_1 \cup \dots \cup I_r}.$$

The group $S_n \times S_d$ acts by permuting the labelling of the markings and scalings. The action of $S_n \times S_d$ on the real locus $\overline{M}^{\mathbb{R}}_{n+1,d}(\mathbb{A})$ has fundamental domain $\overline{M}^{\mathbb{R},\geq 0}_{n,s}(\mathbb{A})$ the closure of the set $M^{\mathbb{R},>0}_{n,s}(\mathbb{A})$ where the markings $z_1 < z_2 < \dots < z_n$ are real, the scalings $\phi_1, \dots, \phi_s$ are real, and $\phi_1 < \phi_2 < \dots < \phi_s$.

Definition 6.3. Let U_0, U_1, U_2 be CohFT's. Given morphisms $\phi_{01} \in \operatorname{Hom}(U_0, U_1)$ and $\phi_{12}) \in \operatorname{Hom}(U_1, U_2)$ define define the *composition of CohFT's* as the sequence of maps

$$(\phi_{12} \circ \phi_{01})^n : U_0^n \otimes H(\overline{M}_{n,2}(\mathbb{A})) \to U_2 \tag{22}$$
$$\sum_{I_1 \cup I_2 \cup \dots \cup I_r = \{1,\dots,n\}} \phi_{12}^r(\phi_{01}^{|I_1|}(\alpha_i, i \in I_1; \cdot), \dots, \phi_{01}^{|I_r|}(\alpha_i, i \in I_r; \cdot); \cdot)(\iota^*_{I_1 \cup \dots \cup I_r}\beta).$$

Here the dots indicate insertion of the Kunneth components of $\iota^*_{I_1 \cup \dots \cup I_r}(\beta)$ with respect to the Kunneth decompositions. The definition of composition of weak morphisms of

CohFT's is similar, except that one allows the subsets $I_1, \ldots, I_r$ in the partition to be empty. A *commutative triangle of CohFT*'s is a triple of morphisms $\phi_{01}, \phi_{12}, \phi_{02}$ such that ϕ_{02} agrees with $\phi_{12} \circ \phi_{01}$ on $U_0^n \otimes H(\overline{M}_{n,2}(\mathbb{A}))$ after composing with the natural restriction map $H(\overline{M}_{n,2}(\mathbb{A})) \to H(\overline{M}_{n,1}(\mathbb{A}))$ defined by restricting to the space where the scalings are equal. Similarly one can define *commutative simplices* of CohFT's of higher dimension.

The need for additional information in the composition of CohFT's seems quite natural, since the composition maps in the CohFT themselves have additional inputs. On the other hand, morphisms of F-manifolds form an ordinary category by composition.

REFERENCES

[1] B. Dubrovin. Geometry of 2D topological field theories. In *Integrable systems and quantum groups (Montecatini Terme, 1993)*, volume 1620 of *Lecture Notes in Math.*, pages 120–348. Springer, Berlin, 1996.

[2] W. Fulton. *Introduction to Toric Varieties*, volume 131 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, 1993.

[3] Ana Rita Pires Gaio and Dietmar A. Salamon. Gromov-Witten invariants of symplectic quotients and adiabatic limits. *J. Symplectic Geom.*, 3(1):55–159, 2005.

[4] Alexander Givental. A mirror theorem for toric complete intersections. In *Topological field theory, primitive forms and related topics (Kyoto, 1996)*, volume 160 of *Progr. Math.*, pages 141–175. Birkhäuser Boston, Boston, MA, 1998.

[5] C. Hertling and Yu. Manin. Weak Frobenius manifolds. *Internat. Math. Res. Notices*, (6):277–286, 1999.

[6] Hiroshi Iritani. Quantum D-modules and generalized mirror transformations. *Topology*, 47(4):225–276, 2008.

[7] Sean Keel. Intersection theory of moduli space of stable n-pointed curves of genus zero. *Trans. Amer. Math. Soc.*, 330(2):545–574, 1992.

[8] Yuri I. Manin. *Frobenius manifolds, quantum cohomology, and moduli spaces*, volume 47 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1999.

[9] S. Ma'u and C. Woodward. Geometric realizations of the multiplihedron and its complexification. 40 pages. arXiv:0802.2120.

[10] Ross Street. Categorical structures. In *Handbook of algebra, Vol. 1*, pages 529–577. North-Holland, Amsterdam, 1996.

[11] C. Woodward and F. Ziltener. Functoriality for Gromov-Witten invariants under symplectic quotients. 2008 preprint.

[12] F. Ziltener. Symplectic vortices on the complex plane. ETH Zurich 2006 Ph.D. thesis.

UNDERGRADUATE PROGRAM, DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139. *E-mail address: nglukhoa@mit.edu*

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854. *E-mail address: ctw@math.rutgers.edu*